Available online at www.sciencedirect.com
SCIENCE DIRECT®
ELSEVIER
Journal of Organometallic Chemistry 669 (2003) 217–218
Journal of Organometallic Chemistry
www.elsevier.com/locate/jorganchem

Contents of Volume 669

Organotransition metal modified sugars: Part 23. Synthesis of vinylcarbene chromium complexes containing a *C*-monosaccharide ligand
E. Janes, K.H. Dötz 1

Addition of diazoalkanes to alkyne(pentacarbonyl)–chromium and –tungsten — formation of 3*H*-pyrazole complexes
M.M. Abd-Elzaher, B. Weibert, H. Fischer 6

Borohydride reduction of a rhenium-bound acetonitrile: an example of a chelating iminoborane ligand at a low valent metal center
A.C. Hillier, T. Fox, H.W. Schmalle, H. Berke 14

Phosphanstabilisierte Silber(I)–Carboxylate: Synthese und Verwendung in der Silber-CVD
H. Schmidt, Y. Shen, M. Leschke, T. Haase, K. Kohse-Höinghaus, H. Lang 25

Synthesis and X-ray structure of palladium dichloride complexed with THF and 2,6,7-trioxa-3,5,8-tris(trichloromethyl)-1,4-diphosphabicyclo[2.2.2]octane
Y. Kim, J.G. Verkade 32

Synthesis and characterization of cyclopentadienyl thiolato complexes of magnesium
A. Xia, M.J. Heeg, C.H. Winter 37

Reproducible high-yield syntheses of $[Ru_3(CO)_{12}]$, $[H_4Ru_4(CO)_{12}]$, and $[Ru_6C(CO)_{16}]^{2-}$ by a convenient two-step methodology involving controlled reduction in ethylene glycol of $RuCl_3 \cdot nH_2O$
E. Lucenti, E. Cariati, C. Dragonetti, D. Roberto 44

Conformationally rigid diphosphine arene–ruthenium(II) complexes as catalysts for transfer hydrogenation of ketones
K.Y. Ghebreyessus, J.H. Nelson 48

Tetranuclear heterodimetallic metallamacrocyles with M–Sn(IV) (M = Mo or W) bonds. Crystal structures of $\{p\text{-}[(CO)_3MoC_5H_4C(O)]_2C_6H_4\}\{(Ph_2Sn)_2S\}$ and $\{p\text{-}[(CO)_3MoC_5H_4C(O)]_2C_6H_4\}\{(Ph_2Sn)_2CH_2\}$
L.-F. Tang, J.-F. Chai, S.-B. Zhao, J.-T. Wang 57

Influence of alkoxy ligands on the Cp–Al bonding mode in $[Cp_2Al\text{-}\mu\text{-}OR]_2$ from X-ray crystallographic and ^{27}Al-NMR spectroscopic solution studies
S. Szumacher, I. Madura, J. Zachara, A.R. Kunicki, A. Cebulski, L. Jerzykiewicz 64

Hydroboration of alkyn-1-yl(methyl)silanes bearing functional substituents at silicon
B. Wrackmeyer, H.E. Maisel, W. Milius, M.H. Bhatti, S. Ali 72

Late transition metal complexes derived from diphosphazane monosulfide ligands: X-ray crystal structures of $[Ru_3(\mu\text{-}CO)(CO)_7(\mu_3\text{-}S)\{Ph_2PN((S)\text{-}^*CHMePh)PPh_2\text{-}\kappa^2P,P\}]$ and $[Rh(CO)Cl\{Ph_2PN((S)\text{-}^*CHMePh)P(S)Ph_2\}\text{-}\kappa^2P,S]$: Part 16. Organometallic chemistry of diphosphazanes
K. Raghuraman, S.S. Krishnamurthy, M. Nethaji 79

The lithiation of arylamines and the preparation of cyclopentadienyltitanium(IV) arylaminate complexes
A.G. Avent, P.B. Hitchcock, G.J. Leigh, M. Togrou 87

Synthesis of titanium(IV) (cyclopentadienylalkyl)aminate complexes
P.B. Hitchcock, G.J. Leigh, M. Togrou 101

Anomalous reaction of an aryl silane with $Co_2(CO)_8$; characterisation of $Me_2NC_6H_4Si[Co(CO)_4][OCCo_3(CO)_9]_2$
G.J. Harfoot, B.K. Nicholson 106

Comparative study of structure–activity relationship of di- and tri-organotin(IV) derivatives of amino acid and peptides
M. Nath, S. Pokharia, G. Eng, X. Song, A. Kumar 109

Elsevier Science B.V.
doi:10.1016/S0022-328X(03)00084-6

Syntheses of a series of trinuclear MIr_2 or pentanuclear MIr_4 bimetallic bis(selenido) and selenido–sulfido clusters (M = Pd, Pt, Fe, Co) from diiridium μ-bis(hydroselenido) and μ-hydroselenido–hydrosulfido complexes $[\{(\eta^5\text{-}C_5Me_5)IrCl\}_2(\mu\text{-}SeH)(\mu\text{-}EH)]$ (E = Se, S)
S. Nagao, H. Seino, M. Hidai, Y. Mizobe 124

Synthesis and reactivity of a new Fe(II) 5-(4-pyridyl)-tetrazolate complex and X-ray structure of its doubly protonated derivative.
A. Palazzi, S. Stagni, S. Selva, M. Monari 135

Preparation and oxidation of polarized Au(III) complexes having both the *C*-deprotonated-2-phenylpyridine (ppy) and a sulfur-rich dithiolate ligand and X-ray crystal structure of $[Au(\eta^2\text{-}C,N\text{-}ppy)(\eta^2\text{-}S,S\text{-}C_8H_4S_8)]\cdot 0.5DMF$
K. Kubo, M. Nakano, H. Tamura, G.-e. Matsubayashi, M. Nakamoto 141

Secondary bonding in *para*-substituted diphenyltellurium dichlorides $(p\text{-}XC_6H_4)_2TeCl_2$ (X = H, Me, MeO) probed by ^{125}Te MAS NMR spectroscopy. Crystal and molecular structure of $(p\text{-}MeC_6H_4)_2TeCl_2$
J. Beckmann, D. Dakternieks, A. Duthie, N.A. Smith 149

Modelling nucleophilic substitution at silicon using solution ^{19}F-NMR chemical shift, $^1J_{Si-F}$ and $^2J_{C-F}$ coupling constant data of pentacoordinate silicon compounds. Correlation with other magnetic nuclei and X-ray structures
A.R. Bassindale, Y.I. Baukov, M. Borbaruah, S.J. Glynn, V.V. Negrebetsky, D.J. Parker, P.G. Taylor, R. Turtle 154

Synthesis, characterisation and study of the first luminescent platinum(II) compound with a $[C,N,S]^-$ terdentate ligand. X-ray crystal structure of $[Pt\{C_6H_4\text{–}CH\text{=}N\text{–}(C_6H_4\text{–}2\text{-}SMe)\}Cl]$
A. Caubet, C. López, X. Solans, M. Font-Bardía 164

Hydridorhodium(I) complexes with amphiphilic polyether phosphines: NMR study and biphasic hydroformylation of 1-octene
A. Solsona, J. Suades, R. Mathieu 172

Synthesis and X-ray structures of some metal carbonyl complexes containing dipyridylformamidine ligands
H.-C. Liang, Y.-Y. Wu, F.-C. Chang, P.-Y. Yang, J.-D. Chen, J.-C. Wang 182

Synthesis, structure, and reactivity of novel iron(II) complexes with a five-membered chelate ligand $\kappa^2(Si,N)$-$SiMe_2O(2\text{-}C_5H_4N)$
T. Sato, M. Okazaki, H. Tobita, H. Ogino 189

The first example of a structurally characterized octahedral hydrotris(pyrazolyl)borate iron methyl complex
O. Graziani, L. Toupet, J.-R. Hamon, M. Tilset 200

Book reviews 207
Erratum 210
Author Index of Volume 669 211
Subject Index of Volume 669 213
Contents Index of Volume 669 217